

Heterotic Modular Invariants and Level–Rank Duality *T. Gannon [‡] and M.A. Walton [†]*Department of Mathematics, York University**Toronto, Ontario, Canada M3J 1P3*

tgannon@mathstat.yorku.ca

*Department of Applied Mathematics and Theoretical Physics**University of Cambridge**Silver Street, Cambridge CB3 9EW, U.K.*

m.a.walton@damtp.cam.ac.uk

Abstract

New heterotic modular invariants are found using the level-rank duality of affine Kac-Moody algebras. They provide strong evidence for the consistency of an infinite list of heterotic Wess-Zumino-Witten (WZW) conformal field theories. We call the basic construction the *dual-flip*, since it flips chirality (exchanges left and right movers) and takes the level-rank dual. We compare the dual-flip to the method of conformal subalgebras, another way of constructing heterotic invariants. To do so, new level-one heterotic invariants are first found; the complete list of a specified subclass of these is obtained. We also prove (under a mild hypothesis) an old conjecture concerning exceptional $A_{r,k}$ invariants and level-rank duality.

PACS: 11.25.Hf, 02.20.Tw

Keywords: conformal field theory, affine algebras

* Supported in part by NSERC.

[‡] Address from July 1998: Math Dept, Univ. Alberta, Edmonton, Canada

[†] On leave from the Physics Dept, Univ. Lethbridge, Alberta, Canada

1. Introduction

Many methods are now known for the direct construction of Wess-Zumino-Witten (WZW) modular invariants. The most important among these start from generic “symmetries” of the modular matrix S .

For example, the WZW models realise affine Kac-Moody algebras as their current algebras. Symmetries of S result from diagram automorphisms of the affine algebras that are not also diagram automorphisms of their horizontal (finite-dimensional) subalgebras. Orbifolds by these symmetries have non-diagonal modular invariants as their torus partition functions [1]. (In the more general context of rational conformal field theory, this orbifold procedure is known as the method of simple currents [2][3].)

The WZW S matrices also have nice arithmetic properties, and so obey certain Galois relations [4]. In [5], these Galois relations were used to construct modular invariants. As for simple currents, one can generalise the Galois considerations to arbitrary rational conformal field theories.

One generic symmetry of WZW modular S matrices sticks out, however. The level-rank duality of WZW models [6][7][8] has not been formulated in a general rational conformal field theory context. More importantly here, it has not been used in a direct manner to construct WZW modular invariants¹. We correct this latter omission.

What we find are new *heterotic* modular invariants, i.e. those describing theories with different holomorphic (left-moving) and anti-holomorphic (right-moving) excitations (see [11][12]). The modular invariants are thus integer (sesquilinear) combinations of characters of one affine algebra with complex conjugates of the characters of a *different* (possibly trivial) algebra. For example, in simple cases, the invariants will be linear combinations of the characters of affine algebras, describing systems – the so-called *meromorphic* conformal field theories – with no right-moving part. In these cases, the construction can be indicated schematically as

$$(g; g) \Rightarrow (g \oplus \tilde{g}; \{\}) \quad . \quad (1.1)$$

Here g indicates an affine nontwisted Kac-Moody algebra $X_{r,k}$, where X_r is the simple Lie horizontal subalgebra of rank r , and k is the fixed level. The semi-colon separates the algebras of the left-moving (L) and right-moving (R) sectors, and $\tilde{g}$ is the algebra that is level-rank dual to g .

¹ Level-rank duality has previously been used in an indirect manner to construct modular invariants [9][10].

To see why, notice that level-rank duality equates the S and T matrices of one WZW model to the *complex conjugates* of the S and T matrices of the dual WZW model (see below), rather than to the dual S and T matrices themselves. To remedy this, we need a way of exchanging the complex conjugates for the dual modular matrices. But this kind of thing has been done in [13], where the modular invariants for certain diagonal coset theories were classified. There a ‘switch’ (or ‘flip’) of left-moving and right-moving fields was used in order to relate the modular transformations of fields of the coset theory to those of the corresponding direct product theory. One of the ways elements of the coset and direct product modular matrices differ is by complex conjugation applied to one factor matrix.

Here we use a switch similar to that of [13] on theories with an affine left-moving current algebra and the same algebra on the right-moving side. Consequently, heterotic invariants are obtained with current algebra plus dual current algebra that are both left-moving (say). In the simplest case then, a purely holomorphic invariant is obtained, as in (1.1). The most general form of the construction is

$$(h_L; h_R) = (g_1 \oplus g_2; g_3 \oplus g_4) \Rightarrow (g_L; g_R) = (g_1 \oplus \tilde{g}_4; g_3 \oplus \tilde{g}_2) \quad . \quad (1.2)$$

Here $h_{L(R)}$ and $g_{L(R)}$ indicate the original and final left(right)-moving algebras, respectively. The symbols g_i can now stand for direct sums of affine nontwisted Kac-Moody algebras.

The key result is that a combination of a level-rank duality transformation and a chirality flip (changing right-moving to left-moving) can map a modular invariant to another, new modular invariant. For short, we will refer to this operation as a *dual-flip*. The lesson seems to be that a level-rank duality transformation is naturally regarded as also exchanging left-movers and right-movers (‘level $\leftrightarrow$ rank’ means ‘Left $\leftrightarrow$ Right’).

The construction does not work as generally as one might think by reading this far, however. Nevertheless, it seems to be a major source of heterotic WZW modular invariants. Since the heterotic modular invariants are generally more difficult to construct than nonheterotic (left-right symmetric) ones [11][12], this is significant progress.

We find meromorphic (i.e. $(g_L; g_R) = (g_L; \{\})$) invariants for

- $g_L = su(n)_m \oplus su(m)_n$ whenever mn is a perfect square congruent to 1 (mod 24);
- $g_L = C_{r,k} \oplus C_{k,r}$ whenever 12 divides rk ;
- $g_L = B_{r,k} \oplus D_{\frac{k}{2}, 2r+1}$ whenever 48 divides $(2r+1)k$;

- $g_L = D_{r,k} \oplus D_{\frac{k}{2},2r}$ whenever k is even and 24 divides rk .

The method of conformal subalgebras [14] provides the only other known general source of heterotic modular invariants. By comparing it with our *dual-flip method* we find more new results. Specifically, we show that some of our invariants can also be derived by conformal embeddings followed by modular invariant contraction [15]. But this is only possible once we have constructed previously unknown *level-one* heterotic physical invariants (recall that only level-one nontwisted affine algebras can have conformal subalgebras).

In the simplest case, these level-one invariants are meromorphic, and so cannot be constructed by the dual-flip. We classify all invariants corresponding to algebras $(X_{r,1}; \{\})$, $(X_{r,1}; X_{s,1})$ and $(X_{r,1} \oplus X_{s,1}; \{\})$. Invariants obtained from these by conformal embeddings, followed by contraction with other invariants, turn out to be of the type derived from the dual-flip. It seems probable that *all* dual-flip invariants can be derived in this fashion, once the appropriate level-one invariant is constructed.

Among the specific examples we find are $c = 24$ meromorphic invariants. These can be related, by conformal embeddings and conjugations, to entries on Schellekens' list [16].

A major result of this paper is our proof of the conjecture of [9] concerning the relation between nonheterotic exceptionals of $h_L = h_R = su(\bar{r})_k$ and $g_L = g_R = su(k)_{\bar{r}}$. There is no bijection between the sets of such physical invariants (unlike the simpler situation for $C_{r,k} \leftrightarrow C_{k,r}$), but under a mild technical condition which we expect to always be satisfied, $su(\bar{r})_k$ will have exceptionals iff $su(k)_{\bar{r}}$ will. This is discussed toward the end of section 5.

We expect that most, if not all, of the physical invariants we find are partition functions of sensible WZW conformal field theories, whether or not they are heterotic. In order to provide further evidence that such theories are consistent, however, one should calculate four-point functions, for example, and verify they satisfy the required properties. Of course, some of the modular invariants we find are the partition functions of known conformal field theories [17].

What we do here, though, is construct new invariants from known ones, many of which are undoubtedly the partition functions of consistent theories. And we believe that since level-rank duality is a duality between the spaces of conformal blocks of theories [18], the dual-flip extends from a map between invariants to a map between theories.

Section 2 treats preliminaries and the construction for the A_r WZW models. In section 3 the results are extended to the other classical simple Lie algebras. Section 4 contains some explicit examples. The full list of heterotic modular invariants for algebras $(X_{r,1}; \{\})$,

$(X_{r,1}; X_{s,1})$ and $(X_{r,1} \oplus X_{s,1}; \{\})$ (for all simple X) is given in section 5, where it is also verified (in certain cases only) that invariants generated from them are of the ‘dual-flip type’. A conclusion is given in section 6.

2. Preliminaries and A_r invariants

Let $X_r^{(1)}$ denote the nontwisted affine Kac-Moody algebra that is the central extension of the loop algebra of the simple Lie algebra X_r . As indicated above, we use $X_{r,k}$ to signify this affine algebra at fixed positive-integer level k . The corresponding set of integrable highest weights has the following set of horizontal projections:

$$P_+(X_{r,k}) = \{ \lambda = \sum_{i=1}^r \lambda_i w^i \quad : \quad \lambda_i \in \mathbf{Z}_{\geq 0}, \sum_{i=1}^r \lambda_i a_i^\vee \leq k \} \quad . \quad (2.1)$$

Here the $a_i^\vee$ are co-marks, and w^i denotes the i -th fundamental weight of X_r . It will often be convenient to introduce the ‘0th affine Dynkin label’ λ_0 defined by

$$\lambda_0 := k - \sum_{i=1}^r \lambda_i a_i^\vee \quad . \quad (2.2)$$

The affine characters $\text{ch}_\lambda(\tau, z, u)$ transform covariantly under the action of the modular group $\text{SL}_2(\mathbf{Z})$. We have

$$\text{ch}_\lambda(\tau + 1, z, u) = \sum_{\mu \in P_+(X_{r,k})} T_\lambda^\mu \text{ch}_\mu(\tau, z, u) = T_\lambda^\lambda \text{ch}_\lambda(\tau, z, u) \quad , \quad (2.3)$$

and

$$\text{ch}_\lambda(-1/\tau, z/\tau, u - z \cdot z/2\tau) = \sum_{\mu \in P_+(X_{r,k})} S_\lambda^\mu \text{ch}_\mu(\tau, z, u) \quad . \quad (2.4)$$

The modular matrices S and T are symmetric and unitary, and have many other interesting properties (see [19], e.g.). For our purposes, the most important relate to the symmetries of the (extended) Coxeter-Dynkin diagram of $X_r^{(1)}$, and to level-rank duality. The diagram symmetries act on weights by permuting their Dynkin labels. There is the involutive charge conjugation C , which is a symmetry of the (unextended) Coxeter-Dynkin diagram of X_r , and so fixes the zeroth node of the $X_r^{(1)}$ diagram. The other important diagram symmetries are often called simple currents, and are related to the centre of the Lie group $\exp(X_r)$ one gets by exponentiating X_r . When the simple current group is generated by a single element, we will denote the generator by J , and any element of that

group by A . The only exception is $X_r = D_r$, with r even. For that case, the two generators will be represented by J and J_s , with the subscript indicating spinor. One finds (see e.g. [20])

$$S_\lambda^{A\mu} = S_\lambda^\mu e^{2\pi i(A^{-1}w^0)\cdot\lambda} = S_\lambda^\mu e^{2\pi i t_A(\lambda)/|A|} , \quad (2.5)$$

where $|A|$ is the order of A , and $t_A(\lambda) := |A| (A^{-1}w^0) \cdot \lambda$ is an integer obtained from the Dynkin labels of the weight λ . The phase in (2.5) is the eigenvalue of an element of the centre of $\exp(X_r)$, with any vector in the representation of highest weight λ as eigenvector.

For example, with $X_{r,k} = A_{r,k}$, we have the diagram symmetry J that permutes the fundamental weights in a cyclic manner, so that

$$J\lambda = J\left(\sum_{i=1}^r \lambda_i w^i\right) = \sum_{i=1}^r \lambda_{i-1} w^i . \quad (2.6)$$

J has order $|J| = r + 1 =: \bar{r}$, and $t_J(\lambda)$ in (2.5) is the $\bar{r}$ -ality

$$t_J(\lambda) = \bar{r}(J^{-1}w^0) \cdot \lambda = \bar{r}w^r \cdot \lambda = \sum_{i=1}^r i\lambda_i . \quad (2.7)$$

It behaves nicely under the action of J :

$$t_J(J^a\lambda) \equiv ka + t_J(\lambda) \pmod{\bar{r}} . \quad (2.8)$$

Using this, and

$$t_{J^a}(\lambda)/|J^a| \equiv t_J(\lambda)/|J| \pmod{1} , \quad (2.9)$$

one can derive a generalisation of (2.5):

$$S_{J^a\lambda}^{J^b\mu} = \exp[2\pi i(bt_J(\lambda) + at_J(\mu) + kab)/\bar{r}] S_\lambda^\mu . \quad (2.10)$$

The action of J on the elements of T is given by

$$T_{J^a\lambda}^{J^a\mu} = \exp[\pi i(-2at_J(\lambda) + ka(\bar{r} - a))/\bar{r}] T_\lambda^\mu . \quad (2.11)$$

Many (and sometimes all) highest weights of $X_{r,k}$, with X_r classical, can be described by Young diagrams. The transpose of a Young diagram corresponds to a highest weight of a different algebra, where the roles of rank and level are interchanged. If λ is the original highest weight, we denote the highest weight corresponding to the transposed diagram by $R\lambda$. The relations between elements of the modular matrices are not so easily described in a uniform, general way. Restricting to $X_{r,k} = A_{r,k}$ then, we find [21] that the map R is not

a bijection between the sets of weights $P_+(A_{r,k})$ and $P_+(A_{k-1,r+1})$. But it is a bijection between the J -orbits of $P_+(A_{r,k})$ and the $\tilde{J}$ -orbits of $P_+(A_{k-1,r+1})$. (Tildes denote objects relevant to the algebra $A_{k-1,r+1} = su(k)_{\bar{r}}$ that is level-rank dual to $A_{r,k} = su(\bar{r})_k$.) More importantly here, one can also construct a bijection R_0 between a subset of $P_+(A_{r,k})$ and a subset of $P_+(A_{k-1,r+1})$. We define the subsets of weights

$$P_+^0(A_{r,k}) := \{ \lambda \in P_+(A_{r,k}) \quad : \quad t_J(\lambda) \equiv 0 \pmod{\bar{r}} \} \quad . \quad (2.12)$$

It can be verified that R_0 given by

$$R_0 \lambda = \tilde{J}^{-t_J(\lambda)/\bar{r}} R \lambda \quad (2.13)$$

is a bijection between $P_+^0(A_{r,k})$ and $\tilde{P}_+^0(A_{k-1,r+1})$. Furthermore [22],

$$S_\lambda^\mu = \sqrt{\frac{k}{\bar{r}}} \tilde{S}_{R_0 \lambda}^{* R_0 \mu} \quad , \quad T_\lambda^\mu T_0^{*0} = \tilde{T}_{R_0 \lambda}^{* R_0 \mu} \tilde{T}_0^{\bar{0}} \quad , \quad (2.14)$$

for all $\lambda, \mu \in P_+^0(A_{r,k})$.

In general, the partition function of a WZW model will take the form of a sesquilinear combination of affine characters:

$$\mathcal{H}(\tau, z, u) = \sum_{\mu \in P_+} \sum_{\nu' \in P'_+} \text{ch}_\mu H_\mu^{\nu'} \text{ch}_{\nu'}^* \quad . \quad (2.15)$$

Notice that we use lower indices for ‘left-movers’ (holomorphic sector) and upper indices for ‘right-movers’ (anti-holomorphic sector). The primes emphasise that the affine algebras may be different on the two ‘sides’ of the theory. For example, P_+ and P'_+ denote the sets of integrable highest weights of the two algebras. If the two affine algebras coincide, the partition function describes a nonheterotic theory. If they do not, it is relevant to a heterotic theory, as in (2.15). If only the left-movers (right-movers) are present, we have a meromorphic (anti-meromorphic) theory. We will use the symbols $\mathcal{H}, H$; $\mathcal{N}, N$; and $\mathcal{M}, M$ to indicate heterotic; nonheterotic; and meromorphic modular invariants, respectively.

By a *physical invariant* we mean a combination of characters of the form (2.15), with $H_\mu^{\nu'} \in \mathbf{Z}_{\geq 0}$ and $H_0^{0'} = 1$, satisfying²

$$\sum_\lambda S_\mu^\lambda H_\lambda^{\nu'} = \sum_{\lambda'} H_\mu^{\lambda'} S_{\lambda'}^{\nu'} \quad , \quad (2.16)$$

² This term is now standard, though somewhat misleading: note that a *physical* invariant might not be realised as the partition function of a consistent theory.

and

$$\sum_{\lambda} T_{\mu}^{\lambda} H_{\lambda}^{\nu'} = \sum_{\lambda'} H_{\mu}^{\lambda'} T_{\lambda'}^{\nu'} . \quad (2.17)$$

In (2.17), the condition $H_0^{0'} = 1$ requires

$$c \equiv c' \pmod{24} , \quad (2.18)$$

which ensures there is no modular anomaly. Here c, c' are the central charges of the two sides of the theory. For example, if the affine algebra on one side is $X_{r,k}$ with X_r simple, then the corresponding central charge is $c(X_{r,k}) = kD/(k+h^{\vee})$, where D is the dimension of X_r , and $h^{\vee}$ is its dual Coxeter number.

It is of some value to study candidate partition functions that satisfy (2.16) exactly, but satisfy (2.17) only when T_{μ}^{λ} and $T_{\lambda'}^{\nu'}$ there are replaced with $T_{\mu}^{\lambda} T_0^{*0}$ and $T_{\lambda'}^{\nu'} T_0'^{*0'}$ respectively (so the modular anomaly condition (2.18) may not be satisfied). We call these objects *anomalous* physical invariants. They can appear as the partition functions of certain sectors of critical string partition functions. For example, the sector of *one* of the factors E_8 in the $E_8 \otimes E_8$ heterotic string is described by such a partition function³. They will also be useful here as simple examples to illustrate our construction.

The following simple observations will be used repeatedly in what follows (proofs can be found in [22]). Let $\mathcal{H}$ be any physical invariant (anomalous or otherwise). Let $\mathcal{J}_{LR}$ be the set of all simple currents $(A; A')$, where A is a simple current of the holomorphic sector, and A' is a simple current of the anti-holomorphic sector, and let $\mathcal{J}(H)$ be the set of all $(A; A') \in \mathcal{J}_{LR}$ such that $H_{A0}^{A'0'} \neq 0$. Then $\mathcal{J}(H)$ is a subgroup of $\mathcal{J}_{LR}$, and for any $(A; A') \in \mathcal{J}_{LR}$,

$$(A; A') \in \mathcal{J}(H) \text{ iff } H_{A\lambda}^{A'\mu'} = H_{\lambda}^{\mu'} \text{ for all } \lambda \in P_+, \mu' \in P'_+ , \quad (2.19)$$

and finally, $(A; A') \in \mathcal{J}(H)$ iff the following selection rule holds:

$$H_{\lambda}^{\mu'} \neq 0 \text{ implies } t_A(\lambda)/|A| \equiv t_{A'}(\mu')/|A'| \pmod{1} \quad (2.20)$$

for all $\lambda \in P_+, \mu' \in P'_+$.

Our objective here is to write heterotic invariants using level-rank duality. In the simplest case, we construct a meromorphic invariant from a known nonheterotic invariant.

³ In heterotic string phenomenology, one E_8 part of the theory can play the role of a *hidden sector* [23]. The possible relevance of heterotic (anomalous) physical invariants to hidden sectors was pointed out by C.S. Lam (as cited in [11]).

Let N_μ^ν and $M_{\mu,\tilde{\nu}}$ denote the coefficient matrices of the original nonheterotic and resulting meromorphic invariant, respectively. In the remainder of this section we will consider only the A_r algebras. Consider first the case where the affine algebra of the invariant $\mathcal{N}$ is $A_{r,k}$ for both left- and right-movers, and the meromorphic invariant $\mathcal{M}$ has algebra $A_{r,k} \oplus A_{k-1,r+1}$. The simplest construction of a meromorphic invariant (we will generalise this later in this section) corresponds to the choice

$$M_{\mu,\tilde{\nu}} = \begin{cases} N_\mu^{R_0\tilde{\nu}} , & \text{for all } \mu \in P_+^0(A_{r,k}), \tilde{\nu} \in \tilde{P}_+^0(A_{k-1,r+1}) , \\ 0, & \text{otherwise} . \end{cases} \quad (2.21)$$

Notice the similarity with the ‘switch’ used in [13]; there left- and right-moving fields were exchanged, while here a right-moving field is traded for a left-mover that is also its level-rank dual. Using the term introduced above, fields are replaced by their flipped-duals.

The construction of (2.21) only works for those physical invariants $\mathcal{N}$ satisfying

$$N_\lambda^\mu \neq 0 \Rightarrow \lambda, \mu \in P_+^0(A_{r,k}) . \quad (2.22)$$

But such invariants exist, and they can also be characterised as obeying $N_{J_0}^0 = N_0^{J_0} = 1$ (2.20). We can find a useful general constraint right away then, by applying T -invariance (2.17), (2.11) to $N_{J_0}^0 = 1$: if r is even, $\bar{r}$ must divide k ; and if r is odd, $2\bar{r}$ must divide k .

To verify modular invariance (2.16), we must show that

$$\sum_{\lambda \in P_+(A_{r,k})} \sum_{\tilde{\zeta} \in P_+(A_{k-1,r+1})} S_\mu^\lambda \tilde{S}_{\tilde{\nu}}^{\tilde{\zeta}} M_{\lambda,\tilde{\zeta}} = M_{\mu,\tilde{\nu}} . \quad (2.23)$$

With our ansatz, the left hand side becomes

$$\sum_{\lambda \in P_+^0(A_{r,k})} \sum_{\tilde{\zeta} \in \tilde{P}_+^0(A_{k-1,r+1})} S_\mu^\lambda N_\lambda^{R_0\tilde{\zeta}} \tilde{S}_{\tilde{\nu}}^{\tilde{\zeta}} , \quad (2.24)$$

where we use (2.22). Since R_0 is a bijection of $P_+^0(A_{r,k})$ onto $\tilde{P}_+^0(A_{k-1,r+1})$, we can write

$$\sum_{\lambda, \zeta \in P_+^0(A_{r,k})} S_\mu^\lambda N_\lambda^\zeta \tilde{S}_{R_0\zeta}^{\tilde{\nu}} = \sum_{\lambda, \zeta \in P_+^0(A_{r,k})} S_\mu^\lambda N_\lambda^\zeta \left(\sqrt{\frac{\bar{r}}{k}} S_{\zeta, R_0\tilde{\nu}}^* \right) , \quad (2.25)$$

using the symmetry of S and (2.14). By the modular invariance of N_μ^ν , this is just $N_\mu^{R_0\tilde{\nu}} \sqrt{\frac{\bar{r}}{k}}$. The S -invariance of $M_{\mu,\tilde{\nu}}$ follows, if we have

$$k = \bar{r} . \quad (2.26)$$

T -invariance is shown in similar fashion, using (2.14). The factors of T_0^{*0} and $\tilde{T}_0^{\tilde{0}}$ are the only complication. But they are absent if the central charge condition (2.18) is satisfied; here it takes the form

$$c(A_{r,k}) + c(A_{k-1,r+1}) = k(r+1) - 1 \equiv 0 \pmod{24} . \quad (2.27)$$

The condition therefore reduces to

$$(r+1)^2 = k^2 \equiv 1 \pmod{24} , \quad (2.28)$$

after imposing (2.26). So the ansatz (2.21) produces a non-anomalous physical invariant iff $k = \bar{r}$ is odd and not divisible by 3; (2.21) is anomalous iff $k = \bar{r}$ is odd and divisible by 3.

There is a way to weaken the stringent requirement (2.26), by generalising (2.21) somewhat. $\bar{r}$ and k must remain odd, but condition (2.26) can be relaxed to

$$\bar{r}k \text{ is a perfect square.} \quad (2.29)$$

To see how, write out the prime decomposition of $\bar{r}/k$ as $\prod_i p_i^{2a_i} \prod_j q_j^{-2b_j}$, where p_i, q_j are distinct primes, and a_i, b_j are positive integers. Define $a := \prod_i p_i^{a_i}$, $b := \prod_j q_j^{b_j}$; then we have $\bar{r}/k = (a/b)^2$, with a and b two coprime integers.

The key is to replace (2.22) with

$$N_\lambda^\mu \neq 0 \Rightarrow t_J(\lambda) \equiv t_J(\mu) \equiv 0 \pmod{\bar{r}/a}. \quad (2.30)$$

Equivalently, we need only demand $N_{J^{a_0}}^0 = N_0^{J^{a_0}} = 1$, by (2.20). Such $\mathcal{N}$ will always exist, for the given choice of a (provided $\bar{r}$ is odd and the perfect-square condition (2.29) is satisfied). (2.21) then gets replaced with

$$M_{J^{ia}\lambda, \tilde{J}^{jb}\tilde{\mu}} = \begin{cases} N_\lambda^{R_0\tilde{\mu}} & , \quad \forall \lambda \in P_+^0(A_{r,k}), \tilde{\mu} \in \tilde{P}_+^0(A_{k-1,r+1}) , \\ 0 & , \quad \text{otherwise} , \end{cases} \quad (2.31)$$

for all i, j . The well-definedness of $\mathcal{N}$ in (2.31), (2.21) and elsewhere is a consequence of (2.19), (2.20).

For (2.31) to be modular invariant, (2.27) and (2.29) must be satisfied. But anomalous invariants will be obtained whenever both k and $\bar{r}$ are odd, 3 divides $k\bar{r}$, and (2.29) is satisfied.

Modular invariance can be verified easily by following the steps used above for (2.21). In this more general case, (2.24) has an extra factor of ab . Then the right hand side of

(2.25) has an extra factor of $ab/a^2 = b/a$, which exactly cancels the $\sqrt{\bar{r}/k}$. So (2.26) gets replaced with the perfect square condition (2.29), and we obtain infinitely many new heterotic invariants for each rank r . That $\bar{r}$ and k must both be odd, follows from T -invariance (2.17) applied to $M_{J^a 0, \tilde{0}} = M_{0, \tilde{J}^b \tilde{0}} = 1$.

The strategy used in constructing the heterotic invariant (2.31) applies in a much more general setting: namely, one where the final holomorphic/antiholomorphic algebras are

$$\begin{aligned} g_L &= su(\bar{r}_1)_{k_1} \oplus su(k_1)_{\bar{r}_1} \oplus \cdots \oplus su(\bar{r}_m)_{k_m} \oplus su(k_m)_{\bar{r}_m} \\ g_R &= su(\bar{s}_1)_{\ell_1} \oplus su(\ell_1)_{\bar{s}_1} \oplus \cdots \oplus su(\bar{s}_n)_{\ell_n} \oplus su(\ell_n)_{\bar{s}_n} \end{aligned} \quad (2.32)$$

where $\bar{s}_i = s_i + 1$. Condition (2.29) will be replaced here by more complicated constraints most conveniently expressed in integral lattice terminology – we won't give the details here.

An example though is the construction of (nonheterotic) physical invariants for $g_L = g_R = su(k)_{\bar{r}}$ from those of $h_L = h_R = su(\bar{r})_k$. Choose any $a|\bar{r}$, $ka^2/\bar{r} \in \mathbf{Z}$, and integers $b|a$ and b' such that $k(b^2 - b'^2)/\bar{r} \in \mathbf{Z}$ (if $\bar{r}$ is even, replace these denominators $\bar{r}$ with $2\bar{r}$). We require $\mathcal{N}$ to be any physical invariant with $(J^a; \text{id}), (\text{id}; J^a), (J^b; J^{b'}) \in \mathcal{J}(N)$, obeying the additional property that for any $\lambda, \mu \in P_+(A_{r,k})$ with $N_\lambda^\mu \neq 0$, there exist simple currents $(A; A') \in \mathcal{J}(N)$ such that $A\lambda, A'\mu \in P_+^0$. This will happen provided $kab/\bar{r}$ is an integer coprime to a . Then we will be able to define a nonheterotic invariant by dual-flipping: choose $\tilde{a}|k$, $\tilde{b}|\tilde{a}$ and $\tilde{b}'$ satisfying $\bar{r}\tilde{a}^2/k, \bar{r}(\tilde{b} - \tilde{b}'^2)/k \in \mathbf{Z}$, and $\bar{r}\tilde{a}\tilde{b}/k$ is an integer coprime to $\tilde{a}$. Then

$$\tilde{N}_{\tilde{J}^{\tilde{a}\ell + \tilde{b}i}\tilde{\lambda}}^{\tilde{J}^{\tilde{a}j + \tilde{b}'i}\tilde{\mu}} = N_{R_0\lambda}^{R_0\mu} \quad (2.33)$$

for any $\lambda \in P_+^0, \mu \in P_+^0$, where all other entries of $\tilde{N}$ vanish. This always works, unless both k and $\bar{r}$ are even and the same power of 2 exactly divides each. If $\mathcal{N}$ is a simple current invariant (i.e. is given by (5.6) below), then so is $\tilde{\mathcal{N}}$. Much more interesting is when $\mathcal{N}$ is exceptional. For example, the level 3 exceptionals of A_r (given in [24]) are dual-shifts of the familiar exceptionals of A_2 , and the level 2 exceptionals (except the $A_{9,2}$ one, which succumbs to the power of 2 condition) are dual-flips of the exceptionals of A_1 . Similar conditions apply to the $A_{3,8} \leftrightarrow A_{7,4}$ and $A_{4,5} \leftrightarrow A_{4,5}$ exceptionals in [22]. It is possible to generalise this correspondence, and indeed we describe at the end of section 5 a generalisation valid for any $\bar{r}, k$ and $\mathcal{N}$ (in particular, it allows us to recover the $A_{9,2}$ exceptional from the $A_{1,10}$ one). This correspondence is clearly of value for the classification of the nonheterotic partition functions of $A_{r,k}$.

3. C_r , B_r and D_r invariants

The other classical algebras can be treated in a straightforward manner using the results of [7].

$X_{r,k} = C_{r,k}$ is simplest. We have an order-two diagram symmetry J that acts on weights in the following way:

$$J\lambda = J\left(\sum_{i=1}^r \lambda_i w^i\right) = \sum_{i=1}^r \lambda_{r-i} w^i . \quad (3.1)$$

The C_r version of (2.5) is

$$S_\lambda^{J\mu} = \exp[\pi i t_J(\lambda)] S_\lambda^\mu , \quad (3.2)$$

with

$$t_J(\lambda) = 2(J^{-1}0) \cdot \lambda / k = 2w^r \cdot \lambda = \sum_{i=1}^r i\lambda_i . \quad (3.3)$$

The action of J on the elements of T is given by

$$T_{J\lambda}^{J\mu} = \exp[\pi i (kr/2 - t_J(\lambda))] T_\lambda^\mu . \quad (3.4)$$

Each weight in $P_+(C_{r,k})$ is in one-to-one correspondence with a Young diagram with no more than r rows, and no more than k columns. For this algebra, the transpose operation R supplies a bijection between the weights of $P_+(C_{r,k})$ and $P_+(C_{k,r})$. The modular matrices obey

$$\tilde{S}_{R\lambda}^{R\mu} = S_\lambda^\mu , \quad \tilde{T}_{R\lambda}^{R\mu} \tilde{T}_{\tilde{0}}^{*\tilde{0}} = T_\lambda^{*\mu} T_0^0 \exp[\pi i t_J(\lambda)] . \quad (3.5)$$

(The $C_{r,k}$ S -matrix is real, hence there is no ‘ $*$ ’ on $\tilde{S}$ in (3.5).) Provided 4 divides rk , the analysis used for $A_{r,k}$ goes through, using

$$M_{\tilde{\mu},\nu} := N_{\tilde{\mu}}^{R_0\nu} , \quad (3.6)$$

where $\mathcal{N}$ is *any* nonheterotic invariant, and $R_0\nu := \tilde{J}^{t_J(\nu)} R\nu$. For a non-anomalous physical invariant, the central charge condition must be satisfied. It becomes

$$c(C_{r,k}) + c(C_{k,r}) = 2kr \equiv 0 \pmod{24} , \quad (3.7)$$

i.e. 12 must divide kr .

The generalisation corresponding to (2.32) is also possible for the C_r algebras. Furthermore, there is no need to stick to one type of algebra. We can start with an invariant with algebras

$$h_L = h_R = g_1 \oplus g_2 \oplus \dots \oplus g_m \oplus h_1 \oplus h_2 \oplus \dots \oplus h_n \quad (3.8)$$

and find one with

$$\begin{aligned} g_L &= g_1 \oplus \tilde{g}_1 \oplus \dots \oplus g_m \oplus \tilde{g}_m \\ g_R &= h_1 \oplus \tilde{h}_1 \oplus \dots \oplus h_n \oplus \tilde{h}_n \quad , \end{aligned} \quad (3.9)$$

with some restrictions. But we will not belabour the point by making this more explicit.

Level-rank duality for the orthogonal algebras, $B_r = so(2r+1)$ and $D_r = so(2r)$, is best treated together [7]. For example, if $so(N)_k$ denotes the affine $so(N)$ algebra at level k , the duality is

$$so(N)_k \leftrightarrow so(k)_N \quad , \quad (3.10)$$

and so does not respect the difference between the B and D algebras.

When they exist, the duality equations for spinor representations of the orthogonal algebras are much more complicated. For this reason, we will restrict ourselves to taking duals of tensor representations only. The highest weights λ of tensor representations of B_r satisfy

$$\lambda_r \equiv 0 \pmod{2} \quad , \quad (3.11)$$

and those of D_r obey

$$\lambda_r \equiv \lambda_{r-1} \pmod{2} \quad . \quad (3.12)$$

Let $P_+^0(so(N)_k)$ denote the corresponding subset of $P_+(so(N)_k)$. In both cases the condition can be written as

$$t_J(\lambda) := w^1 \cdot \lambda \equiv 0 \pmod{1} \quad , \quad (3.13)$$

where J is the following simple current:

$$\begin{aligned} J\lambda &= \lambda_0 w^1 + \lambda_2 w^2 + \dots + \lambda_r w^r \quad \text{for } B_r \quad , \\ J\lambda &= \lambda_0 w^1 + \lambda_2 w^2 + \dots + \lambda_{r-2} w^{r-2} + \lambda_r w^{r-1} + \lambda_{r-1} w^r \quad \text{for } D_r \quad . \end{aligned} \quad (3.14)$$

There is one remaining complication for the case of D_r . The irreducible tensor representations of $so(N)$ are in one-to-one correspondence with Young diagrams having the sum of the lengths of their first two columns less than or equal to N , except when the first column is of length $N/2$. If λ denotes a weight of D_r , then let $C\lambda$ denote the weight obtained by exchanging the Dynkin labels λ_r and λ_{r-1} . (For convenience define $C\lambda = \lambda$

for weights of B_r .)⁴ A diagram with first column length $N/2 = r$ corresponds to the pair $[\lambda] := \{\lambda, C\lambda\}$. Again the level-rank dual R is given by the transpose operation on Young diagrams. So it will be more convenient to consider level-rank duality for the C -orbits $[\lambda]$. For this purpose, define

$$S_{[\lambda]}^{[\mu]} := \sum_{\substack{\kappa \in [\lambda] \\ \nu \in [\mu]}} S_{\kappa}^{\nu} , \quad T_{[\lambda]}^{[\mu]} := \sum_{\kappa \in [\lambda]} T_{\kappa}^{\mu} . \quad (3.15)$$

First, let us write the level-rank duality relation for T [7]:

$$T_{[\lambda]}^{[\lambda]} T_{[0]}^{*[0]} = \exp[\pi i t'(\lambda)] \tilde{T}_{R[\lambda]}^{*R[\lambda]} \tilde{T}_{[\tilde{0}]}^{[\tilde{0}]} , \quad (3.16)$$

valid for all $so(N)_k$, with

$$t'(\lambda) = \sum_{i=1}^{r-1} i\lambda_i + r\lambda_r/2 , \quad \text{for } so(N)_k = B_{r,k} , \quad (3.17)$$

and

$$t'(\lambda) = \sum_{i=1}^{r-2} i\lambda_i + \frac{r-2}{2}\lambda_{r-1} + \frac{r}{2}\lambda_r , \quad \text{for } so(N)_k = D_{r,k} . \quad (3.18)$$

Notice that for a tensor representation of highest weight λ , $t'(\lambda) \in \mathbf{Z}$. Also, for $D_{r,k}$, $t'(\lambda) \equiv t_{J_s}(\lambda)$ or $\frac{1}{2}t_{J_s}(\lambda) \pmod{2}$, for r even or odd resp., where J_s is the simple current given by

$$\begin{aligned} J_s \lambda &= \sum_{i=1}^r \lambda_{r-i} w^i \quad \text{for } r \text{ even} , \\ J_s \lambda &= \lambda_r w^1 + \sum_{i=2}^r \lambda_{r-i} w^i \quad \text{for } r \text{ odd} . \end{aligned} \quad (3.19)$$

The level-rank duality relation for S is [7]:

$$S_{[\lambda]}^{[\mu]} = \tilde{S}_{R[\lambda]}^{R[\mu]} , \quad \forall \lambda, \mu \in P_+^0(so(N)_k) . \quad (3.20)$$

Incidentally, the level-rank duality $so(N)_k \leftrightarrow so(k)_N$ extends in the natural way to small N , using $so(3)_k \cong A_{1,2k}$, $so(4)_k \cong A_{1,k} \oplus A_{1,k}$, $so(5)_k \cong C_{2,k}$, and $so(6)_k \cong A_{3,k}$.

Consider first $h_L = h_R = B_{r,k}$. We require the nonzero entries N_{λ}^{μ} of the initial invariant to obey $\lambda, \mu \in P_+^0(B_{r,k})$, which means

$$N_{J_0}^0 = N_0^{J_0} = 1 . \quad (3.21)$$

⁴ As defined here, C is not the charge conjugation for D_r with r even; it is for the other cases.

This forces k to be even, in which case such invariants will always exist. When 16 divides k , we can get a $B_{r,k} \oplus D_{k/2, 2r+1}$ meromorphic invariant by putting

$$M_{\lambda, J_s^j C^i R_0 \mu} := N_{\lambda}^{\mu} , \quad (3.22)$$

where $R_0 \lambda := J^{t'(\lambda)} R \lambda$, and $i, j \in \{0, 1\}$ (the simple current $J^{t'}$ cancels the sign picked up in (3.16), but forces the presence of the J_s^j , which is what requires the constraint $16|k$). (Another way of cancelling the sign in (3.16) would be to use instead $J_s^{t'}$, but again this requires $16|k$ and yields nothing new: $M\tilde{C}$ instead of M .)

The other alternative is $h_L = h_R = D_{r,k}$ with k even. As before, we require the nonzero entries N_{λ}^{μ} to satisfy $\lambda, \mu \in P_+^0(D_{r,k})$. Also, the fact that (3.20) only holds at the level of C -orbits, means that C should commute with N . Thus we obtain the constraints

$$N_{J_0}^0 = N_0^{J_0} = 1 \text{ and } N_{\lambda}^{\mu} = N_{C\lambda}^{C\mu} \quad \forall \lambda, \mu . \quad (3.23)$$

Such invariants will always exist, since k is even.

Now to cancel the sign in (3.16), we must use $J_s^{t'}$, which requires 8 to divide rk . The formula becomes

$$M_{\lambda, J^j C^i R^0 \mu} := N_{\lambda}^{\mu} , \quad (3.24)$$

where $R^0 \lambda := J_s^{t'(\lambda)} R \lambda$ or $J_s^{2t'(\lambda)} R \lambda$, for r even or odd resp., and $i, j \in \{0, 1\}$.

In both cases, the constraint on N, k is that 16 divides Nk . For a non-anomalous physical invariant, we need only impose the central charge condition

$$c(so(N)_k) + c(so(k)_N) = \frac{Nk}{2} \equiv 0 \pmod{24} , \quad (3.25)$$

or 48 must divide Nk .

4. Examples

In order to write some simple examples of the new physical invariants in a concise way, we introduce a little notation. Let $[\lambda_0, \lambda_1, \dots, \lambda_r]$ represent the *character* ch_{λ} , with $\lambda = \sum_{i=1}^r \lambda_i w^i$, where we put $\lambda_0 := k - \sum_{i=1}^r \lambda_i a_i^{\vee}$ as above. So, the trivial diagonal physical invariant for algebra $X_{r,k}$ would be written as

$$\sum_{\lambda \in P_+(X_{r,k})} |[\lambda_0, \lambda_1, \dots, \lambda_r]|^2 \quad (4.1)$$

in this notation.

We will content ourselves here with simple examples involving the A_r and C_r algebras only.

A large class of nonheterotic invariants for $X_{r,k}$, with X_r simple, is the class of partition functions of orbifolds by subgroups of the centre of $\exp(X_r)$ [20][25]. Let us first discuss meromorphic invariants that can be constructed from them.

A manifestly non-negative formula was written in [26] for these invariants:

$$N(\exp(X_r)/Z_n)_\mu^\lambda = \sum_{m=0}^{n-1} \delta_\mu^{A^m \lambda} \delta_1 \{ (Aw^0) \cdot \left(\lambda + \frac{1}{2}mkAw^0 \right) \} . \quad (4.2)$$

It is valid for any cyclic orbifold group Z_n , as long as

$$\frac{nk(Aw^0) \cdot (Aw^0)}{2} \equiv 0 \pmod{1} . \quad (4.3)$$

Here A generates the corresponding diagram symmetry subgroup, so that $|A| = n$, and

$$\delta_y \{x\} = \begin{cases} 1 & , \text{ if } x/y \text{ is integer;} \\ 0 & , \text{ otherwise.} \end{cases} \quad (4.4)$$

First consider nonheterotic $A_{r,k}$ physical invariants $\mathcal{N} = \sum_{\lambda,\mu} \text{ch}_\lambda N_\lambda^\mu \text{ch}_\mu^*$ of this orbifold type. For $A_{r,k}$, the outer automorphism in (4.2) can be $A = J^d$ for any d dividing $\bar{r}$. However, in order to construct the simplest meromorphic physical invariants $\mathcal{M} = \sum_{\lambda,\mu} \text{ch}_\lambda \text{ch}_\mu M_{\lambda,\mu}$, i.e. using (2.21), we have seen in (2.22),(2.26) that we must put $A = J$, and also take $k = \bar{r}$ odd. (We will consider the more general (2.31) later.)

First, we will ignore the central charge condition (2.28), and find a simple anomalous meromorphic physical invariant. The lowest value of $k = \bar{r}$ at which a non-diagonal invariant (4.2) exists is 3. Eqn. (4.2) yields

$$\mathcal{N} = |[3, 0, 0] + [0, 3, 0] + [0, 0, 3]|^2 + 3 |[1, 1, 1]|^2 , \quad (4.5)$$

and (2.21) produces

$$\mathcal{M}^{(an.)} = ([3, 0, 0] + [0, 3, 0] + [0, 0, 3])^2 + 3 ([1, 1, 1])^2 . \quad (4.6)$$

For a non-anomalous physical invariant, (2.28) requires $k^2 \equiv 1 \pmod{24}$. But this just means that $k = \bar{r}$ must be coprime to 6, so that the construction works for all even r that

are not congruent to 2 (mod 3). An *infinite series* of meromorphic physical invariants is thus obtained. The smallest possible rank is $r = 4$, where

$$\begin{aligned} \mathcal{N} = & |J\{5, 0, 0, 0, 0\}|^2 + |J\{3, 1, 0, 0, 1\}|^2 + |J\{3, 0, 1, 1, 0\}|^2 \\ & + |J\{1, 2, 0, 0, 2\}|^2 + |J\{1, 0, 2, 2, 0\}|^2 + 5|[1, 1, 1, 1, 1]|^2, \end{aligned} \quad (4.7)$$

where we have defined

$$A\{\lambda_0, \lambda_1, \dots, \lambda_r\} := \sum_{a=1}^{|A|} A^a [\mu_0, \mu_1, \dots, \mu_r]. \quad (4.8)$$

In general, R is a bijection between the set of J -orbits in $P_+(A_{r,k})$ and the set of $\tilde{J}$ -orbits in $P_+(A_{k-1,r+1})$. R_0 is a bijection between $P_+^0(A_{r,k})$ and $\tilde{P}_+^0(A_{k-1,r+1})$. If $D := \gcd\{\bar{r}, k\}$, $P_+^0(A_{r,k})$ and $\tilde{P}_+^0(A_{k-1,r+1})$ are composed of complete $J^{\bar{r}/D}$ -orbits and $\tilde{J}^{k/D}$ -orbits, respectively. So, when $\bar{r} = k$, R_0 maps the J -orbits of $P_+^0(A_{r,k})$ onto the $\tilde{J}$ -orbits of $\tilde{P}_+^0(A_{k-1,r+1})$. Consequently, the meromorphic $(A_{4,5} \oplus A_{4,5}; \{\})$ invariant that results from (4.7) is

$$\begin{aligned} \mathcal{M} = & J\{5, 0, 0, 0, 0\}^2 + J\{3, 1, 0, 0, 1\}^2 + J\{3, 0, 1, 1, 0\} J\{1, 2, 0, 0, 2\} \\ & + J\{3, 0, 1, 1, 0\} J\{1, 2, 0, 0, 2\} + J\{1, 0, 2, 2, 0\}^2 + 5[1, 1, 1, 1, 1]^2, \end{aligned} \quad (4.9)$$

which corresponds to the 67th entry in the table of $c = 24$ meromorphic theories given in [16].

The invariants for larger $\bar{r} = k$ are too long to write here, so we will instead write a general, formal equation. Let $P_{+,J}(A_{r,\bar{r}})$ denote a set of representatives of the J -orbits in $P_+^0(A_{r,\bar{r}})$. Then the general form of the nonheterotic $A_{r,\bar{r}}$ physical invariant is

$$\mathcal{N} = \sum_{\lambda \in P_{+,J}(A_{r,\bar{r}})} \frac{\bar{r}}{\|\langle \lambda \rangle_J\|} |\langle \text{ch}_\lambda \rangle_J|^2, \quad (4.10)$$

with $\langle \lambda \rangle_J$ being the J -orbit of λ . By (2.21), we construct the following meromorphic physical invariant

$$\mathcal{M} = \sum_{\lambda \in P_{+,J}(A_{r,\bar{r}})} \frac{\bar{r}}{\|\langle \lambda \rangle_J\|} (\langle \text{ch}_\lambda \rangle_J) \left(\langle \tilde{\text{ch}}_{R_0 \lambda} \rangle_{\tilde{J}} \right), \quad (4.11)$$

as long as (2.28) is satisfied.

An example of an (anomalous) invariant of the type (2.31) is also too long to write here, but a general equation similar to (4.10) can easily be found.

So far we have only considered examples of meromorphic invariants that can be obtained from the nonheterotic orbifold invariants (4.2). Thus we have not yet considered any example of an exceptional invariant. There is no obstruction to using such invariants, however. Consider the exceptional invariant of $(A_{4,5}; A_{4,5})$ [22]:

$$\begin{aligned} \mathcal{N} = & |J\{0, 0, 5, 0, 0\}|^2 + J\{0, 1, 3, 1, 0\} [1^5]^* + [1^5] J\{0, 1, 3, 1, 0\}^* + \\ & |J\{1, 0, 3, 0, 1\}|^2 + |J\{0, 2, 1, 2, 0\}|^2 + |J\{2, 0, 1, 0, 2\}|^2 + 4 |[1^5]|^2 . \end{aligned} \quad (4.12)$$

The invariant obtained by (2.21) is then just

$$\begin{aligned} \mathcal{M} = & (J\{0, 0, 5, 0, 0\})^2 + J\{0, 1, 3, 1, 0\} [1^5] + [1^4] J\{0, 1, 3, 1, 0\} + \\ & J\{1, 0, 3, 0, 1\} J\{0, 2, 1, 2, 0\} + J\{0, 2, 1, 2, 0\} J\{1, 0, 3, 0, 1\} + \\ & (J\{2, 0, 1, 0, 2\})^2 + 4 ([1^5])^2 . \end{aligned} \quad (4.13)$$

This is the ninth entry in the table of $c = 24$ meromorphic theories given in [16].

Next we treat $C_{r,k}$, where 4 divides rk . Any physical invariant $\mathcal{N}$ will generate a (possibly anomalous) meromorphic invariant. Generically, there will be two physical invariants $\mathcal{N}$ for such r, k : they correspond to the choices $A = id$ and $A = J$ in (4.2). The resulting $\mathcal{M}$ will be anomalous unless the central charge condition (3.7) is satisfied, i.e. unless 12 divides kr .

Four simple anomalous heterotic invariants can be found for $kr = 4$. With $r = k = 2$, (4.2) gives the $(C_{2,2}; C_{2,2})$ invariant

$$\mathcal{N} = |[2, 0, 0] + [0, 0, 2]|^2 + 2 |[0, 2, 0]|^2 + 2 |[1, 0, 1]|^2 . \quad (4.14)$$

From this, the anomalous $(C_{2,2} \oplus C_{2,2}; \{\})$ invariant

$$\mathcal{M}^{(an.)} = ([2, 0, 0] + [0, 0, 2])^2 + 2[0, 2, 0][1, 0, 1] + 2[1, 0, 1][0, 2, 0] \quad (4.15)$$

is obtained. For $\mathcal{N}$ the diagonal invariant of $C_{2,2}$, we get

$$\begin{aligned} \mathcal{M}^{(an.)} = & [2, 0, 0]^2 + [0, 0, 2]^2 + [1, 1, 0][0, 1, 1] + [0, 1, 1][1, 1, 0] \\ & + [1, 0, 1][0, 2, 0] + [0, 2, 0][1, 0, 1]. \end{aligned} \quad (4.16)$$

With $r = 4, k = 1$, we find

$$\mathcal{N} = |[1, 0, 0, 0, 0] + [0, 0, 0, 0, 1]|^2 + 2 |[0, 0, 1, 0, 0]|^2 , \quad (4.17)$$

and the resulting anomalous $(C_{4,1} \oplus A_{1,4}; \{\})$ invariant

$$\mathcal{M}^{(an.)} = ([1, 0, 0, 0, 0] + [0, 0, 0, 0, 1])([4, 0] + [0, 4]) + 2[0, 0, 1, 0, 0][2, 2] . \quad (4.18)$$

Choosing $\mathcal{N}$ diagonal instead gives us

$$\mathcal{M}^{(an.)} = [1, 0^4][4, 0] + [0, 1, 0^3][1, 3] + [0, 0, 1, 0, 0][2, 2] + [0^3, 1, 0][3, 1] + [0^4, 1][0, 4] . \quad (4.19)$$

The lowest value of kr at which non-anomalous physical invariants can be written is 12. As an example, we choose $r = 4, k = 3$. Eqn. (4.2) with $A = J$ yields

$$\begin{aligned} \mathcal{N} = & |J\{3, 0, 0, 0, 0\}|^2 + 2|[0, 0, 3, 0, 0]|^2 \\ & + |J\{2, 0, 1, 0, 0\}|^2 + |J\{2, 0, 0, 0, 1\}|^2 \\ & + |J\{1, 2, 0, 0, 0\}|^2 + |J\{0, 2, 1, 0, 0\}|^2 \\ & + |J\{1, 0, 0, 2, 0\}|^2 + |J\{1, 0, 2, 0, 0\}|^2 \\ & + |J\{1, 1, 0, 1, 0\}|^2 + 2|[0, 1, 1, 1, 0]|^2 \\ & + 2|[1, 0, 1, 0, 1]|^2 . \end{aligned} \quad (4.20)$$

The meromorphic $C_{4,3} \oplus C_{3,4}$ physical invariant we obtain from it is

$$\begin{aligned} \mathcal{M} = & J\{3, 0, 0, 0, 0\} \tilde{J}\{4, 0, 0, 0\} + 2[0, 0, 3, 0, 0][2, 0, 0, 2] \\ & + J\{2, 0, 1, 0, 0\} \tilde{J}\{2, 2, 0, 0\} + J\{2, 0, 0, 0, 1\} \tilde{J}\{0, 4, 0, 0\} \\ & + J\{1, 2, 0, 0, 0\} \tilde{J}\{3, 0, 1, 0\} + J\{0, 2, 1, 0, 0\} \tilde{J}\{2, 1, 0, 1\} \\ & + J\{1, 0, 0, 2, 0\} \tilde{J}\{1, 0, 3, 0\} + J\{1, 0, 2, 0, 0\} \tilde{J}\{2, 0, 2, 0\} \\ & + J\{1, 1, 0, 1, 0\} \tilde{J}\{1, 2, 1, 0\} + 2[0, 1, 1, 1, 0][1, 1, 1, 1] \\ & + 2[1, 0, 1, 0, 1][0, 2, 2, 0] , \end{aligned} \quad (4.21)$$

which corresponds to the 70th entry in [16].

Notice that the algebra $C_{4,3} \oplus C_{3,4}$ does not satisfy a condition of [16]: that the ratio of the dual Coxeter number to the level be equal for all $X_{r,k}$ in the affine algebra. But that is because the condition is derived assuming that the affine algebra is maximal. Clearly, it is satisfied by the $D_{24,1}$ algebra in which $C_{4,3} \oplus C_{3,4}$ is conformally embedded. We will explain how (4.21) can be recovered from this embedding at the end of the next section.

5. Level-one holomorphic physical invariants and progeny

There is an intimate connection between level-rank dualities and certain conformal embeddings. For example, the conformal subalgebra $su(\bar{r})_k \oplus su(k)_{\bar{r}} \subset su(\bar{r}k)_1$ contains much information about the duality $su(\bar{r})_k \leftrightarrow su(k)_{\bar{r}}$. So it is natural to suspect that some of the invariants we find by dual-flipping may also be obtainable (less directly) by the method of conformal subalgebras.

But the simplest of our invariants are meromorphic. To derive them using conformal subalgebras, we would need to start with meromorphic level-one physical invariants for X_r , i.e. physical invariants for $(X_{r,1}; \{\})$. Previous to this work, only a few such invariants were known – for some of these, see e.g. [16].

Motivated by this situation, however, we were led to derive such invariants. Furthermore, we were able to classify all *heterotic* invariants for algebras $(X_{r,1}; \{\})$, $(X_{r,1}; X_{s,1})$ and $(X_{r,1} \oplus X_{s,1}; \{\})$. The complete list for $(X_{r,1}; \{\})$ follows, using the notation $\text{ch}_i := \text{ch}_{w^i}$.

(a) $(A_{r,1}; \{\})$, $\bar{r} = s^2, \bar{r}$ odd : (anomalous unless 24 divides r)

$$\mathcal{M} = \text{ch}_0 + \text{ch}_s + \text{ch}_{2s} + \dots + \text{ch}_{(s-1)s} .$$

(d) $(D_{r,1}; \{\})$, $r \equiv 0 \pmod{8}$: (anomalous unless 24 divides r)

$$\mathcal{M}_1 = \text{ch}_0 + \text{ch}_r ,$$

$$\mathcal{M}_2 = \text{ch}_0 + \text{ch}_{r-1} .$$

$\text{ch}_r, \text{ch}_{r-1}$ are the highest weights of the spinor and conjugate-spinor representations of $D_{r,1}$, respectively.

(e8) $(E_{8,1}; \{\})$: $\mathcal{M} = \text{ch}_0$ is anomalous.

(*) $(X_{r,1}; \{\})$ for $X_r = B_r, C_r, E_6, E_7, F_4, G_2$: no invariants.

The completeness of the above list follows immediately from the simple level-one nonheterotic (i.e. $(X_{r,1}; X_{r,1})$) classification [27]: if $\sum \text{ch}_i$ is a heterotic invariant (possibly anomalous), then $|\sum \text{ch}_i|^2$ is a nonheterotic physical invariant. So we can simply run down the list of [27], to see if any of those invariants can be written as a perfect square.

We will quickly run through the classification for $(X_{r,1} \oplus X_{s,1}; \{\})$ and $(X_{r,1}; X_{s,1})$. Its proof is also straightforward; T -invariance takes one almost all the way.

(a1) $(A_{r,1}; A_{s,1})$ has invariants provided $\sqrt{rs} \in \mathbf{Z}$ and $r \equiv s \pmod{2}$; they will be anomalous unless $r \equiv s \pmod{24}$.

(a2) $(A_{r,1} \oplus A_{s,1}; \{\})$ has invariants provided $\sqrt{rs} \in \mathbf{Z}$, both $\bar{r}$ and $\bar{s}$ are odd, and every prime $p \equiv 3 \pmod{4}$ divides $\bar{r}$ exactly an even number of times; they will be anomalous unless $r \equiv -s \pmod{24}$.

(b1) $(B_{r,1} \oplus B_{s,1}; \{\})$ has exactly 1 invariant, provided $r + s \equiv 7 \pmod{8}$; it will be anomalous unless $r + s \equiv -1 \pmod{24}$.

(b2) $(B_{r,1}; B_{s,1})$ has exactly 1 invariant, provided $r \equiv s \pmod{8}$; it will be anomalous unless $r \equiv s \pmod{24}$.

(d1) $(D_{r,1} \oplus D_{s,1}; \{\})$ will have no invariants, unless $r \equiv -s \pmod{8}$; it will have exactly 6 if either $r \equiv s \equiv 0$ or $r \equiv s \equiv 4 \pmod{8}$, otherwise it will have exactly 2 provided $r \equiv -s \pmod{8}$; all of these will be anomalous unless $r \equiv -s \pmod{24}$.

(d2) $(D_{r,1}; D_{s,1})$ will have no invariants, unless $r \equiv s \pmod{8}$; it will have exactly 6 if either $r \equiv s \equiv 0$ or $r \equiv s \equiv 4 \pmod{8}$, otherwise it will have exactly 2 (provided $r \equiv s \pmod{8}$); all of these will be anomalous unless $r \equiv s \pmod{24}$.

The only other heterotic invariant⁵ of type $(X_{r,1} \oplus X_{s,1}; \{\})$ or $(X_{r,1}; X_{s,1})$ for X simple, is the anomalous one $\mathcal{M} = \text{ch}_0 \text{ch}_0$ for $X_r = X_s = E_8$.

The invariants in (b1), (b2), (d1) and (d2) are easy to write down. Those in (a1) and (a2) are much messier and should be interpreted in the language of self-dual lattices.

Some of the invariants found above seem to be partition functions for strings on heterotic nonsimply-connected group manifolds. That is, as partition functions they appear to describe closed bosonic strings with left-moving sector and right-moving sector propagating on *different* nonsimply-connected group manifolds. Perhaps the simplest example is (a), where the group would be $SU(s^2)/Z_s$, and the string meromorphic.

We want to show that invariants found in the previous section by dual-flipping can also be found using conformal subalgebras and contraction. The appropriate conformal subalgebras are what we dub the *level-rank* conformal subalgebras, for obvious reasons: the prototype is $su(m)_n \oplus su(n)_m \subset su(mn)_1$. In this work, we will consider only this level-rank conformal subalgebra.

First let us write down its conformal branching rules [9][21]. Let us again use the notation of the last section, where a weight denotes its corresponding affine character. Then we have

$$w^j = \sum_{\substack{\lambda \in P_+(A_{m-1}, n) : \\ t_J(\lambda) \equiv j \pmod{m}}} \lambda(\tilde{J}^{[j-t_J(\lambda)]/m} R\lambda) \quad , \quad (5.1)$$

⁵ For the choice $X = C$, we have been able to show this only by assuming the invariant is non-anomalous, i.e. obeys (2.18), although we expect there will be no anomalous heterotics either.

where here w^j denotes the character of the $su(mn)_1$ representation restricted to the subalgebra. This last formula is consistent under the swap of $su(m)_n$ and $su(n)_m$, as must be. That's because $\tilde{t}_J(R\lambda) \equiv t_J(\lambda) \pmod{n}$ and (2.8) imply

$$\tilde{t}_J(\tilde{J}^{[j-t_J(\lambda)]/m} R\lambda) \equiv j \pmod{n} . \quad (5.2)$$

With this result we will be able to show that the level-one invariant of (a) above yields the meromorphic invariants (2.21) and (2.31) for $\mathcal{N}$ given by (4.2) with $A = J$ and $A = J^a$, respectively. Substituting (5.1), with $m = n = s$, into the $(su(s^2)_1; \{\})$ invariant $\sum_{a=0}^{s-1} w^{as}$ yields

$$\sum_{a=0}^{s-1} \sum_{\lambda \in P_+^0(A_{s-1,s})} \lambda \left(J^{a-t_J(\lambda)/s} R\lambda \right) = \sum_{\lambda \in P_+^0(A_{s-1,s})} \lambda \left(\sum_{a=0}^{s-1} J^a R_0 \lambda \right) , \quad (5.3)$$

using (2.13). But this equals (4.11) with $\bar{r} = s$. More generally, when $\bar{r}k = s^2$, this construction yields (2.31) for $\mathcal{N}$ given by the choice $A = J^a$ in (4.2).

To recover other invariants, such as (4.13), we must also use the technique of modular invariant contraction [15]: that is, contracting one modular invariant with a second produces another modular invariant. For example, let $M'_{\lambda,\nu}$ be the coefficient matrix of the meromorphic $(A_{4,5} \oplus A_{4,5}; \{\})$ invariant (4.9), a special case of (5.3), just obtained by conformal embedding. Then, if N'_λ is the matrix for the exceptional $(A_{4,5}; A_{4,5})$ invariant (4.12), the contraction

$$\frac{1}{5} \sum_{\nu \in P_+(A_{4,5})} M'_{\lambda,\nu} N'_\nu = M_{\lambda,\mu} \quad (5.4)$$

yields the matrix elements of the exceptional $(A_{4,5} \oplus A_{4,5}; \{\})$ invariant (4.13). The normalisation $\frac{1}{5}$ in (5.4) is needed to get $M_{0,0} = 1$; in general this necessary step could introduce fractions into the modular invariant. The possibility of fractions is the main difficulty with any contraction formula.

It appears that any invariant obtainable by the dual-flip can also be recovered from a level-one invariant by the combination of level-rank conformal subalgebras and modular invariant contraction. We have found no exception to this rule. For example, consider the large set of $(su(k)_{\bar{r}}; su(k)_{\bar{r}})$ invariants described as the dual-flips of $(su(\bar{r})_k; su(\bar{r})_k)$ invariants by (2.33). We will follow the procedure given in [9]. Start with the diagonal $(su(k\bar{r})_1; su(k\bar{r})_1)$ invariant, and apply the conformal embedding $su(\bar{r})_k \oplus su(k)_{\bar{r}} \subset su(k\bar{r})_1$, to get an invariant $\mathcal{N}''$ for $(su(\bar{r})_k \oplus su(k)_{\bar{r}}; su(\bar{r})_k \oplus su(k)_{\bar{r}})$. Let $\mathcal{N}$ be any

$(su(\bar{r})_k; su(\bar{r})_k)$ physical invariant, and let ℓ be the smallest number $1 \leq \ell \leq \bar{r}$ for which $N_{J_0^\ell}^{J_0} \neq 0$. Define the contraction

$$\tilde{N}_\mu^{\mu'} = \frac{\ell}{\bar{r}} \sum_{\lambda, \lambda' \in P_+(A_{r,k})} N_{\lambda, \mu}^{\lambda', \mu'} N_{\lambda'}^\lambda. \quad (5.5)$$

It turns out (using (2.19),(2.20)) that with the normalisation $\frac{\ell}{\bar{r}}$, $\tilde{\mathcal{N}}$ will be a *physical invariant* for $(su(k)_{\bar{r}}; su(k)_{\bar{r}})$ (this could only be conjectured in [9]).

However, *there is no bijection between the $(su(k)_{\bar{r}}; su(k)_{\bar{r}})$ and $(su(\bar{r})_k; su(\bar{r})_k)$ physical invariants*. This is already evident in [24]. What we obtain here is just as important. It is one of the main results of the paper, and so we will state it more precisely. We call a physical invariant $\mathcal{N}$ a *simple current invariant* if it obeys the selection rule

$$N_\lambda^\mu \neq 0 \Rightarrow \lambda = A\mu \text{ for some simple current } A \quad (5.6)$$

(A depends on λ, μ). We will call a physical invariant $\mathcal{N}$ *exceptional* if neither $\mathcal{N}$ nor any conjugation $C_1 \mathcal{N} C_2$ is a simple current invariant (for A_r , the only nontrivial conjugation is the charge conjugation C , defined by $(C\lambda)_i = \lambda_{\bar{r}-i}$).

Fact (a) If $\mathcal{N}$ is a physical invariant for $(su(\bar{r})_k; su(\bar{r})_k)$, then $\tilde{\mathcal{N}}$ defined by (5.5) will be a physical invariant for $(su(k)_{\bar{r}}; su(k)_{\bar{r}})$.

(b) Suppose $su(\bar{r})_k$ has an exceptional invariant $\mathcal{N}$ obeying $N_{J_0}^{J_0} \neq 0$. Then $su(k)_{\bar{r}}$ will have an exceptional invariant obeying $\tilde{N}_{J_0}^{\tilde{J}_0} \neq 0$.

More generally, if $N_\lambda^{\lambda'} \neq 0$, $t_J(\lambda) \equiv t_J(\lambda') \pmod{\bar{r}}$, and $\lambda' \neq J^i \lambda$ for any i (respectively, $\lambda' \neq C^i J^j \lambda$ for any i, j), then $\tilde{\mathcal{N}}$ will have the corresponding property. This should be of considerable value in future attempts at $A_{r,k}$ physical invariant classifications. Such weights λ, λ' can be found for any known exceptional invariant of $A_{r,k}$. So we can confidently predict the following (which we know to be true if either $\bar{r}$ or k is a product of distinct odd primes):

Conjecture $(su(\bar{r})_k; su(\bar{r})_k)$ will have an exceptional physical invariant, iff $(su(k)_{\bar{r}}; su(k)_{\bar{r}})$ does.

On the other hand, there *is* a bijection between the nonheterotic physical invariants of $C_{r,k}$, and those of $C_{k,r}$, given by the formula $N_\lambda^\mu = \tilde{N}_{R\lambda}^{R\mu}$.

As promised, we conclude this section with a comment on the $c=24$ invariant (4.21). As stated previously, it corresponds in Schellekens' table [16] to the 70th entry, with algebra

$D_{24,1}$. The two invariants of (d) above, with $r=24$, differ by a conjugation (not charge conjugation) and also correspond to entry 70. The invariant (4.21) can be obtained from one of them by the conformal embedding $C_{4,3} \oplus C_{3,4} \subset D_{24,1}$, with branching rules given in [10]. The other (d) invariant produces a different $(C_{4,3} \oplus C_{3,4}; \{\})$ invariant. But it too can be derived by the dual-flip, this time from the diagonal non-heterotic invariant of either $C_{4,3}$ or $C_{3,4}$.

6. Conclusion

To summarise, we have shown how many new heterotic physical invariant combinations of affine Kac-Moody characters can be found. The new invariants are intimately related with the level-rank duality of affine Kac-Moody algebras. They can be derived either of two ways: (i) by the new dual-flip method (see e.g. (2.21)), or (ii) by applying together the two old methods of conformal embeddings [14] and modular invariant contraction [15] (by nonheterotic invariants).

The advantage of the first method is that it is the most direct and explicit. To use the old method (ii), it was first necessary to find new level-one heterotic invariants by force. Method (i) also shows explicitly that level-rank duality is responsible for the existence of the new invariants. This is satisfying, since other generic symmetries of the modular matrices, such as Galois [5] and simple-current relations [2][3], have been shown to produce invariants.

Level-rank duality appears in method (ii) when those conformal subalgebras related to it (such as $A_{m-1,n} \oplus A_{n-1,m} \subset A_{mn-1,1}$) are used. But this points out an advantage of the second method: conformal subalgebras besides the ‘level-rank’ ones can be used. In this sense, method (ii) is more general. We have not investigated this generalisation here, except for the following remark. The heterotic invariants in (a) and (d) above, together with conformal embeddings such as [28] $A_{n,n-1} \subset A_{(n-1)(n+2)/2,1}$, $B_{n,2} \subset A_{2n,1}$, $A_{2n,2n+1} \subset D_{2n(n+1),1}$, $B_{2n,4n-1} \subset D_{n(4n+1)}$, $C_{2n,2n+1} \subset D_{n(4n+1),1}$, $D_{2n,4n-2} \subset D_{n(4n-1),1}$, give us new meromorphic invariants of type $(X_{r,k}; \{\})$ (e.g. for $A_{49,48}$ and $A_{16,17}$).

We should emphasise, however, that compared to nonheterotic invariants, heterotic invariants are rare, as the results of [11][12] show. In contrast, the dual-flip seems to produce a large class of heterotic invariants. So, although we made no attempt at completeness in this paper, it may still be that this method can produce most heterotic invariants for the algebras with level-rank duality. A natural question then is, have we exhausted all generic ways of constructing heterotics?

This question is not premature, considering what seems to be the main (and unexpected) lesson being learned from the nonheterotic physical invariant classifications: the ‘obvious’ ways to construct these physical invariants (most notably, conjugations, simple currents, conformal embeddings, and rank-level duality) succeed in constructing almost all of them.

A direct challenge to this optimism is provided by the meromorphic CFTs, when 8 divides c . There is at least one of these associated with each even self-dual lattice of dimension c . In dimension 24, there are only 24 such lattices, but in dimension 32 there are at least 8 million, and this lower bound grows quickly with the dimension. So there will be an enormous number of $c = 48$ meromorphic CFTs, very few of which can be constructed using these “generic” methods. However, these meromorphic CFTs will in general *not* be of the WZW type considered in this paper. Our point is merely that if one restricts to an affine algebra, and varies the level, then what one finds is very few if any truly exceptional physical invariants. The intractability of an explicit classification of all meromorphic CFTs, which is inherited from the intractability of an explicit classification of all even self-dual lattices, has led many (prematurely, in our opinion) to regard the WZW physical invariant classification as hopeless.

Acknowledgements

MW thanks Matthias Gaberdiel for helpful conversations and the High Energy Physics group of DAMTP for hospitality.

References

- [1] D. Gepner and E. Witten, *Nucl. Phys.* **B278** (1986) 493
- [2] A.N. Schellekens and S. Yankielowicz, *Nucl. Phys.* **B327** (1989) 673
- [3] K. Intriligator, *Nucl. Phys.* **B332** (1990) 541
- [4] A. Coste and T. Gannon, *Phys. Lett.* **B323** (1994) 316
- [5] J. Fuchs, B. Gato-Rivera, B. Schellekens and C. Schweigert, *Phys. Lett.* **B334** (1994) 113
- [6] A. Kuniba and T. Nakanishi, in:
S. Das et al. (editors), *Modern Quantum Field Theory* (World Scientific, New York, 1991) (proceedings of the International Colloquium on Modern Quantum Field Theory, Bombay, 1990)
- [7] E. Mlawer, S. Naculich, H. Riggs and H. Schnitzer, *Nucl. Phys.* **B352** (1991) 863
- [8] T. Nakanishi and A. Tsuchiya, *Comm. Math. Phys.* **144** (1992) 351
- [9] M.A. Walton, *Nucl. Phys.* **B322** (1989) 775
- [10] D. Verstegen, *Comm. Math. Phys.* **137** (1991) 567
- [11] T. Gannon, *Nucl. Phys.* **B402** (1993) 729
- [12] T. Gannon and Q. Ho-Kim, *Nucl. Phys.* **B425** (1994) 319
- [13] T. Gannon and M.A. Walton, *Comm. Math. Phys.* **173** (1995) 175
- [14] A. Bais and A. Taormina, *Phys. Lett.* **B181** (1986) 87
- [15] P. Bouwknegt, *Nucl. Phys.* **B290** (1987) 507
- [16] A.N. Schellekens, *Comm. Math. Phys.* **153** (1993) 159
- [17] I. Dolan, P. Goddard and P. Montague, *Phys. Lett.* **B236** (1990) 165
- [18] S. Naculich and H. Schnitzer, *Nucl. Phys.* **B347** (1990) 687;
T. Nakanishi and A. Tsuchiya, *Comm. Math. Phys.* **144** (1992) 351
- [19] T. Gannon and M.A. Walton, *On fusion algebras and modular matrices*, q-alg/9709039 (1997)
- [20] D. Bernard, *Nucl. Phys.* **B288** (1987) 628
- [21] D. Altschuler, M. Bauer and C. Itzykson, *Comm. Math. Phys.* **132** (1990) 349
- [22] T. Gannon, *Kac-Peterson, Perron-Frobenius, and the classification of conformal field theories*, q-alg/9510026 (1995)
- [23] P. Candelas, G. Horowitz, A. Strominger, and E. Witten, *Nucl. Phys.* **B258** (1985) 46
- [24] T. Gannon, *Lett. Math. Phys.* **39** (1997) 289
- [25] D. Altschuler, J. Lacki and P. Zaugg, *Phys. Lett.* **B205** (1988) 281
G. Felder, K. Gawedzki and A. Kupiainen, *Comm. Math. Phys.* **117** (1988) 127

- [26] C. Ahn and M.A. Walton, *Phys. Lett.* **B223** (1989) 343
- [27] C. Itzykson, *Nucl. Phys. Proc. Suppl.* **5B** (1988) 150
T. Gannon, *Nucl. Phys.* **B396** (1993) 708
- [28] F.A. Bais and P.G. Bouwknecht, *Nucl. Phys.* **B279** (1987) 561;
A.N. Schellekens and N. Warner, *Phys. Rev.* **D34** (1986) 3092